

RP00614

Leader in Biomolecular Solutions for Life Science



Recombinant Human SIRP-gamma/CD172g Protein

Catalog No.: RP00614 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	55423	Q9P1W8

Tags

C-Fc

Synonyms

SIRPG;CD172g;SIRP-B2;SIRPB2;SIRPgamma;bA77C3.1

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Signal-Regulatory Protein Gamma (SIRPG) is a member of the signal-regulatory protein (SIRP) family and also belongs to the immunoglobulin superfamily. SIRPG is detected in the liver, and at very low levels in the brain, heart, lung, pancreas, kidney, placenta, and skeletal muscle. SIRPG is an immunoglobulin-like cell surface receptor. On binding with CD47, SIRPG mediates cell-cell adhesion. Engagement on T-cells by CD47 on antigen-presenting cells results in enhanced antigen-specific T-cell proliferation and costimulates T-cell activation. SIRPG as receptor-type transmembrane glycoproteins is involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes.

Basic Information

Description


Recombinant Human SIRP-gamma/CD172g Protein is produced by Human cells expression system. The target protein is expressed with sequence (Glu29-Pro360) of human SIRP-gamma/CD172g (Accession #Q9P1W8) fused with an Fc tag at the C-terminus.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human SIRP-gamma/SIRPB2/CD172g Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.