ABclonal®

Recombinant Human uPAR/PLAUR Protein

Catalog No.: RP00656 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 5329 Q03405-1

Tags C-His&Avi

Synonyms

U-PAR; uPAR; CD87; PLAUR; MO3; UPAR

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by SDS-PAGE;≥ 95 % as determined by

HPLC.

Endotoxin

 $< 1 EU/\mu g$ of the protein by LAL method.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com

Background

The receptor (u-PAR) for urokinase plasminogen activator (u-PA) is a three-domain protein, GPI-anchored to the cell surface, which focuses the enzymatic activity of u-PA, and allows the cell surface activation of plasminogen. Regulation of the activity of u-PA is also mediated by u-PAR.

Basic Information

Description

Recombinant Human uPAR/PLAUR Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu23-Gly305) of Human uPAR/PLAUR (Accession #Q03405-1) fused with a C-His&Avi tag at the C-terminus.

Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.


Validation Data



Recombinant Human uPAR/PLAUR Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Human uPAR is greater than 95% as determined by SEC-HPLC.



Immobilized Human uPAR, His Tag at 2 μ g/mL (100 μ L/well) on the plate. Dose response curve for Biotinylated Human Human PLAU, His Tag with the EC₅₀ of 22.5ng/mL determined by ELISA.