# Recombinant Human LR3-IGF-1(E51R) Protein

Catalog No.: RP00825 Recombinant

# Sequence Information

Species	Gene ID	Swiss Prot
<i>E.</i>	3479	P05019
coli		

Tags

NO-Tag

### Synonyms

IGF1; IBP1; IGF-1; Insulin-like growth factor 1; Insulin-like growth factor I; IGF-I; Mechano growth factor; MGF; Somatomedin-C

# **Product Information**

Source	Purification
<i>E. coli</i>	≥ 95 % as
	determined by
	SDS-PAGE

ed by

### Endotoxin

< 1 EU/µg of the protein by LAL method.

### Formulation

Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

# Contact

www.abclonal.com G

## Background

Long R3 IGF-1 (LR3 IGF-1) is a synthetic analog of IGF-1 that is generated by modifying the AA sequence for mature human IGF-1. These modifications include the substitution of an Arg for Glu at position three of the mature IGF-1 sequence and the addition of a thirteen aa N-terminal extension, which is derived from methionyl porcine Growth Hormone. These aa changes generate a protein that is still capable of binding to IGF-1 and Insulin receptors, but shows considerably lower affinity binding to IGFBPs compared to wild-type IGF-1. As a result, LR3 IGF-1 has an increased half-life and displays increased biological potency compared to IGF-1.

# **Basic Information**

### Description

Recombinant Human LR3-IGF-1(E51R) Protein is produced by E. coli expression system. The target protein is expressed with sequence (Gly49-Ala118) of human IGF-1 (Accession #P05019) with the substitution of an Arg for Glu at position three and the addition of a thirteen aa N-terminal extension.

### **Bio-Activity**

1、Measured in a cell proliferation assay using FDC-P1 cells. The ED<sub>50</sub> for this effect is 9.85-39.4 ng/mL, corresponding to a specific activity of 2.54×10<sup>4</sup>~1.01×10<sup>5</sup> units/mg.

### Storage


Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.



# Validation Data



Measured in a cell proliferation assay using FDC-P1 cells. The  $ED_{50}$  for this effect is 9.85-39.4 ng/mL, corresponding to a specific activity of  $2.54 \times 10^4 \sim 1.01 \times 10^5$  units/mg.