Recombinant Mouse Flt3 ligand/Flt3L Protein

Catalog No.: RP01058 Recombinant 1 Publications

Sequence Information

Background

Species Gene ID Swiss Prot HEK293 cells 14256 P49772

Tags C-hFc&His

Synonyms

FLT3LG;FL;FLT3L;Flt3 ligand;FLT3LG

Product Information

Purification Source HEK293 cells > 85% by SDS-PAGE.

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

Ð

www.abclonal.com

Basic Information

Description

Recombinant Mouse Flt3 ligand/Flt3L Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Met1-Arg188) of mouse Flt-3 Ligand (Accession #NP_038548.3.) fused with an Fc, 6×His tag at the Cterminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Mouse FLT3, His Tag at 3 µg/mL (100 µL/well) can bind Recombinant Mouse Flt-3 Ligand, The EC₅₀ of Mouse Flt-3 Ligand is 36 ng/mL.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.




Validation Data

170kDa — 130kDa —	
100kDa —	
70kDa —	_
55kDa —	
40kDa —	
35kDa —	
25kDa —	
15kDa —	
10kDa —	


Recombinant Mouse Flt3 ligand/Flt3L Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Recombinant Mouse FLT3, His Tag at 3μ g/mL (100 μ L/well) can bind Recombinant Mouse Flt-3 Ligand, The EC₅₀ of Mouse Flt-3 Ligand is 36 ng/mL.



Immobilized recombinant Human FLT3 at 0.5μ g/mL (100 μ L/well) can bind FLT3L with a linear range of 0.05-1.3 ng/mL.



Recombinant Mouse Flt3 ligand/Flt3L Protein promote the proliferation using Human OCI-AML5 cells. The ED_{50} for this effect is 1.3-5.2ng/mL, corresponding to a specific activity of $1.92 \times 10^5 \sim 7.69 \times 10^5$ units/mg.