

# **Recombinant Human Mesothelin/MSLN Protein**

Catalog No.: RP01220 Recombinant

## **Sequence Information**

**Species Gene ID Swiss Prot** HEK293 cells 10232 Q13421-3

Tags

C-hFc&His

**Synonyms** 

MPF; SMRP; MSLN; SMRP

### **Product Information**

Source

**Purification** 

HEK293 cells

≥ 95 % as determined by SDS-PAGE.

## **Endotoxin**

 $< 0.1 \; \text{EU/}\mu\text{g}$  of the protein by LAL method.

## **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

## Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

### Contact

•

www.abclonal.com

# Background

## **Basic Information**

### Description

Recombinant Human Mesothelin/MSLN Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu296-Gly580) of human Mesothelin (Accession #NP\_005814.2) fused with a Fc, 6×His tag at the C-terminus.

### **Bio-Activity**

Loaded Human Mesothelin/MSLN Protein, C-hFc&His (Catalog: RP01220) on ProA Biosensor, can bind Human CA125 Protein, C-His&Avi (Catalog: RP01417) with an affinity constant of 17.2 nM as determined in BLI assay (Gator).

#### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

<sup>\*</sup> For your safety and health, please wear a lab coat and disposable gloves when handling.


## **Validation Data**



Recombinant Human
Mesothelin/MSLN Protein was
determined by SDS-PAGE under
reducing conditions with Coomassie
Rlue



Immobilized Human CA125 at 5  $\mu$ g/mL (100  $\mu$ L/well) can bind Human Mesothelin/MSLN with a linear range of 0.3-17 ng/mL.



Loaded Human Mesothelin/MSLN Protein, C-hFc&His (Catalog: RP01220) on ProA Biosensor, can bind Human CA125 Protein, C-His&Avi (Catalog: RP01417) with an affinity constant of 17.2 nM as determined in BLI assay (Gator).