

RP01584

Leader in Biomolecular Solutions for Life Science



Recombinant Mouse Hepatocyte growth factor/HGF Protein

Catalog No.: RP01584 **Recombinant**

Sequence Information

Species HEK293 cells
Gene ID 15234
Swiss Prot Q08048

Tags

C-His

Synonyms

SF; NK1; NK2; HGF/SF; SF/HGF; C230052L06Rik;HGF

Product Information

| Source | Purification |
|--------------|---|
| HEK293 cells | ≥ 95 % as determined by SDS-PAGE; ≥ 95 % as determined by HPLC. |

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

HGF, also known as Scatter Factor and Hepatopoietin A, is a pleiotropic protein in the Plasminogen subfamily of S1 peptidases. It is a multidomain molecule that includes an N-terminal PAN/APPLE-like domain, four Kringle domains, and a serine proteinase-like domain that has no detectable protease activity. Mouse HGF is secreted as an inactive 728 amino acid (aa) single chain propeptide. It is cleaved after the fourth Kringle domain by a serine protease to form bioactive disulfide-linked HGF with a 60kDa alpha and 30 kDa beta chain. Alternate splicing generates an isoform that lacks the peptidase and the second, third, and fourth Kringle domains. Mouse HGF shares 91%-95%aa sequence identity with bovine, canine, feline, human, and rat HGF. HGF binds heparan-sulfate proteoglycans and the widely expressed receptor tyrosine kinase, HGF R/c-MET. HGF dependent c-MET activation is implicated in the development of many human cancers. HGF regulates epithelial morphogenesis by inducing cell scattering and branching tubulogenesis. HGF induces the up regulation of integrin alpha 2 beta 1 in epithelial cells by a selective increase in alpha 2 gene transcription. This integrin serves as a collagen I receptor, and its blockade disrupts epithelial cell branching tubulogenesis. HGF can also alter epithelium morphology by the induction of nectin-1 alpha ectodomain shedding, an adhesion protein component of adherens junctions. In the thyroid, HGF induces the proliferation, motility, and loss of differentiation markers of thyrocytes and inhibits TSH-stimulated iodine uptake. HGF promotes the motility of cardiac stem cells in damaged myocardium.

Basic Information

Description

Recombinant Mouse Hepatocyte growth factor/HGF Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln33-Leu728) of mouse Hepatocyte growth factor/HGF (Accession #NP_001276387.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured in a cell proliferation assay using Mv.1.lu cells. The ED₅₀ for this effect is 1.2-5.0 ng/mL, corresponding to a specific activity of 2×10⁵ ~ 8.3×10⁵ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



The purity of Recombinant Mouse Hepatocyte growth factor/HGF Protein is greater than 95% as determined by SEC-HPLC.