# ABclonal www.abclonal.com

## **Recombinant Human Periostin/PN/POSTN Protein**

Catalog No.: RP01748 Recombinant

## **Sequence Information**

**Species Gene ID Swiss Prot** HEK293 cells 10631 Q15063

**Tags** C-6His

Synonyms

PN; OSF2; OSF-2; PDLPOSTN; POSTN

## **Product Information**

**Source** Purification HEK293 cells

**Endotoxin** 

< 1EU/µg

#### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

## Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## **Contact**

€

www.abclonal.com

## **Background**

Periostin ( POSTN ), also known as OSF2 (osteoblast specific factor 2), is a heterofunctional secreted extracellular matrix (ECM) protein comprised of four fasciclin domains that promotes cellular adhesion and movement, as well as collagen fibrillogenesis. Postn is frequently overexpressed in various types of human cancers, stimulating metastatic growth by promoting cancer cell survival, invasion and angiogenesis, and can be a useful marker to predict the behavior of cancer.

### **Basic Information**

## Description

Recombinant Human Periostin/PN/POSTN Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asn22-Gln836) of human Periostin/PN/POSTN (Accession #NP\_006466.2) fused with a 6×His tag at the C-terminus.

#### **Bio-Activity**

Measured by its ability to induce adhesion of ATDC5 mouse chondrogenic cells. Immobilized Recombinant Human Periostin/OSF $\square$  2 at 10  $\mu$ g/mL (100  $\mu$ L/well) induces >19% cell adhesion.

#### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

<sup>\*</sup> For your safety and health, please wear a lab coat and disposable gloves when handling.

## **Validation Data**



Recombinant human Periostin/PN/POSTN Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 100-110 kDa.