

RP01798

Leader in Biomolecular Solutions for Life Science



Recombinant Human WFDC2/HE4/WAP5 Protein

Catalog No.: RP01798

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	10406	Q14508

Tags

C-hFc

Synonyms

HE4; WAP5; EDDM4;
dJ461P17.6; VEGFR-3; FLT-4; WFDC2

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.01EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

WAP four-disulfide core domain protein 2, also known as Epididymal secretory protein E4, Major epididymis-specific protein E4, Putative protease inhibitor WAP5, WFDC2 and HE4, is a secreted protein that contains two WAP domains. WFDC2 / HE4 is a member of a family of stable 4-disulfide core proteins that are secreted at high levels. It is expressed in a number of normal tissues, including male reproductive system, regions of the respiratory tract and nasopharynx. It is highly expressed in a number of tumors cells lines, such ovarian, colon, breast, lung and renal cells lines. Initially described as being exclusively transcribed in the epididymis. WFDC2 may be a component of the innate immune defences of the lung, nasal and oral cavities and suggest that WFDC2 functions in concert with related WAP domain containing proteins in epithelial host defence. WFDC2 re-expression in lung carcinomas may prove to be associated with tumour type and should be studied in further detail. Mammary gland expression of tammar WFDC2 during the course of lactation showed WFDC2 was elevated during pregnancy, reduced in early lactation and absent in mid-late lactation. WFDC2 / HE4 can undergo a complex series of alternative splicing events that can potentially yield five distinct WAP domain containing protein isoforms.

Basic Information

Description

Recombinant Human WFDC2/HE4/WAP5 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu31-Phe124) of human WFDC2/HE4/WAP5 (Accession #NP_006094.3) fused with hFc tag at the C-terminus.


Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Human WFDC2/HE4/WAP5 Protein was resolved with SDS PAGE under reducing (R) and non-reducing (NR) conditions, showing single bands at 45-50 kDa and 75-100 kDa, respectively.