ABclonal www.abclonal.com

Recombinant Human ICOS/CD278 Protein

Catalog No.: RP02255 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 29851 Q9Y6W8

Tags

C-mFc&His

Synonyms

AILIM; CD278; CVID1;ICOS;CD278;CVID1

Product Information

Purification Source HEK293 cells

> 95% by SDS-

PAGE.

Endotoxin

Formulation

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 33 % trehalose is added as protectants before lyophilization.

Reconstitution

Reconstitute with deionized water

Background

The protein encoded by this gene belongs to the CD28 and CTLA-4 cell-surface receptor family. It forms homodimers and plays an important role in cell-cell signaling, immune responses, and regulation of cell proliferation.

Basic Information

Description

Recombinant Human ICOS/CD278 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (ICOS(Glu21-Lys140)&mFc(Pro99-Lys330)) of human ICOS (Accession #) fused with a Fc and His Tag at the C-terminal. .

Bio-Activity

ELISA plate pre-coated by 2 μg/mL (100 μL/well) Human ICOS, mFc-His tagged protein can bind Anti-ICOS Neutralizing antibody in a linear range of 0.24-31.25 ng/mL.

Store the lyophilized protein at -20°C to -80°C for long term.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up


Avoid repeated freeze/thaw cycles.

Contact


0

www.abclonal.com

Validation Data



Recombinant Human ICOS/CD278 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 41.1 kDa



ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human ICOS, mFc-His tagged protein can bind Anti-ICOS Neutralizing antibody in a linear range of 0.24-31.25ng/ml.