Biotinylated Recombinant Human Prolyl endopeptidase FAP Protein

Catalog No.: RP02352 Recombinant

Sequence Information

Background

Basic Information

Description

SpeciesGene IDSwiss ProtHEK293 cells 2191Q12884

Tags

N-His&Avi

Synonyms

DPPIV; FAPA; FAPalpha; SIMP;FAP; DPPIV; prolyl endopeptidase FAP;FAPA;FAPalpha;SIMP, Fibroblast activation protein alpha;FAP

Product Information

Source Purification HEK293 cells > 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com

Bio-Activity Immobilized Biotinylated Human FAP,His Tag at 2 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Anti-FAP Antibody with the EC₅₀ of 17.6 ng/mL

FAP fused with His tag and Avi tag at the N-terminal.

Biotinylated Recombinant Human FAP Protein is produced by Expi293 expression

system. The target protein is expressed with sequence (Leu26-Asp760) of Human

Storage

determined by ELISA.

Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

www.abclonal.com

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Biotinylated Human FAP on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



The purity of Biotinylated Human FAP is greater than 95% as determined by SEC-HPLC.



Immobilized Anti-FAP Antibody at 5μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Human FAP, His Tag with the EC₅₀ of 0.14 μ g/ml determined by ELISA.