Recombinant Cynomolgus PVRIG Protein

Catalog No.: RP02609 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells 102127364		A0A2K5WVV
		8-1

Tags C-hFc

Synonyms

C7orf15; CD112R; MGC104322; MGC138297; MGC2463; PVRIG; ALS2CR18; ALS2CR9; LPD; PREL-2; PREL2; RalGDS/AF-6; RMO1

Product Information

Source	Purification	Storage
HEK293 cells	> 95% as determined by	Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year fro the date of receipt.
	Tris-Bis PAGE [°] ; > 95% as determined	After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-for up to 1 week.
	by HPLC	Avoid repeated freeze/thaw cycles.
Fudataviu		

Endotoxin

Less than 1EU per μ g by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com G

Background

PVRIG (poliovirus receptor related immunoglobulin domain-containing protein), also known as CD112 receptor (CD112R), is an approximately 34 kDa single transmembrane protein in the poliovirus receptor-like protein (PVR) family.PVRIG is a cell surface receptor for NECTIN2. May act as a coinhibitory receptor that suppresses T-cell receptor-mediated signals. Following interaction with NECTIN2, inhibits T-cell proliferation. Competes with CD226 for NECTIN2-binding.

Basic Information

Description

Recombinant Cynomolgus PVRIG Protein is expressed from Expi293 with hFc tag at the C-terminal. ; It contains Thr41-Asp171.

Bio-Activity

* For your safety and health, please wear a lab coat and disposable gloves when handling.



Validation Data



Cynomolgus PVRIG on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Cynomolgus PVRIG is greater than 95% as determined by SEC-HPLC.



Cynomolgus PVRIG, hFc Tag captured on CM5 Chip via Protein A can bind Human Nectin-2, His Tag with an affinity constant of 0.23 μ M as determined in SPR assay (Biacore T200).



Immobilized Cynomolgus PVRIG, hFc Tag at 0.1μ g/ml (100μ l/well) on the plate. Dose response curve for Biotinylated Anti-PVRIG Antibody, hFc Tag with the EC₅₀ of 5.4ng/ml determined by ELISA.