

RP02670

Leader in Biomolecular Solutions for Life Science



Recombinant Human CRTAM/CD355 (A65V) Protein

Catalog No.: RP02670

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	56253	O95727-1

Tags

C-His

Synonyms

CD355 antigen; CD355; CRTAM

Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE.

Endotoxin

< 0.01 EU/μg of the protein by LAL method

Formulation

Reconstitution

Centrifuge the tube before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Class-I Restricted T Cell-Associated Molecule (CRTAM) is a protein that is expressed after T cell activation. The interaction of CRTAM with its ligand, nectin-like 2 (Nect2), is required for the efficient production of IL-17, IL-22, and IFNγ by murine CD4 T cells, and it plays a role in optimal CD8 T and NK cell cytotoxicity. CRTAM promotes the pro-inflammatory cytokine profile; therefore, it may take part in the immunopathology of autoimmune diseases such as diabetes type 1 or colitis.

Basic Information

Description


Recombinant Human CRTAM Protein is expressed from HEK293 cells with a His tag at the C-terminal. ; It contains Ser18-Gly287.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CRTAM/CD355 (A65V) Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized Human CRTAM, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-CRTAM Antibody, hFc Tag with the EC₅₀ of 57.3ng/ml determined by ELISA.