ABclonal www.abclonal.com

Recombinant Mouse FSTL3 Protein

Catalog No.: RP03254 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 83554 Q9EQC7

Tags

C-His

Synonyms

Follistatin-like protein 3; FLRG; Follistatin-like 3; FSTL3

Product Information

Source Purification
HEK293 cells > 95% by Tris-

> 95% by Tris-Bis PAGE and HPLC.

Endotoxin

< 1.0 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com

Background

Follistatin-like 3 (FSTL3) is a novel cytokine that regulates insulin sensitivity and counteracts activin/myostatin signalling. In the present study, regulation of FSTL3 in renal dysfunction was investigated in both human chronic kidney disease (CKD) and acute kidney dysfunction (AKD). Furthermore, mFSTL3 expression was analysed in insulin-sensitive tissues in a mouse model of CKD.

Basic Information

Description

Recombinant Mouse FSTL3 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val24-Val256) of Mouse FSTL3 (Accession #Q9EQC7) fused with His tag at the C-terminus.


Bio-Activity

1. Immobilized Human Activin A, No Tag at 0.2 μ g/mL (100 μ L/well) on the plate. Dose response curve for Mouse FSTL3, His Tag with the EC₅₀ of 18.0 ng/mL determined by ELISA. | 2. Mouse FSTL3, His Tag captured on CM5 Chip via anti-his antibody can bind Human Activin A, No tag with an affinity constant of 53.09 pM as determined in SPR assay (Biacore T200). | 3. Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED₅₀ for this effect is typically 5-25 ng/mL in the presence of 7.5 ng/mL rhActivin A.


Storage

Store at -20° C. Store the lyophilized protein at -20° C to -80° C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20° C for 3 months, at $2-8^{\circ}$ C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Mouse FSTL3 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 35-40 kDa.



The purity of Mouse FSTL3 is greater than 95% as determined by SEC-HPLC.



Immobilized Human Activin A, No Tag at $0.2 \mu g/mL$ ($100 \mu L/well$) on the plate. Dose response curve for Mouse FSTL3, His Tag with the EC₅₀ of 18.0 ng/mL determined by ELISA.